



Danish in-kind simulation efforts to the ESS - an overview

Klinkby, Esben Bryndt; Willendrup, Peter Kjær; Lefmann, Kim

Publication date:
2012

[Link back to DTU Orbit](#)

Citation (APA):

Klinkby, E. B. (Author), Willendrup, P. K. (Author), & Lefmann, K. (Author). (2012). Danish in-kind simulation efforts to the ESS - an overview. Sound/Visual production (digital) <http://www.nonboel.dk/nordic-gen4/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Danish in-kind simulation efforts to the ESS


- an overview

Esben Klinkby, DTU Nutech

contributions from:

Peter Willendrup (DTU)

Kim Lefmann (Uni Copenhagen)



Outline

- From hard protons to cold neutrons
 - Experimental overview
 - Simulation tools
 - “*In house*” simulation efforts
- “*In country*” instrument design

Risø




Target station design

- Target station modeled with MCNPX
- Spallation takes place in rotating tungsten target
- The scale of the objects under study dictates the use of cold neutrons ($\sim 1-10\text{meV}$)
- Neutrons are moderated in H_2 and H_2O for cold and thermal neutrons




EUROPEAN
SPALLATION
SOURCE



Flip-view




EUROPEAN
SPALLATION
SOURCE




Neutron guides

- Ni and Ti are chemically similar, but have very different refraction indices
- Coating with alternating layers: “Supermirrors” of which guides can be built which ~without loss, transport cold neutrons to radiation safe distances allowing ToF to be measured





Collimators & slits

- Works from the principle: Absorb anything which don't have the desired direction (gadolinium)
- Discriminate beam in *space* and *divergence*



Disk Choppers

- Introduce pulses
- Discriminate beam in *time*
- Combining two choppers \sim Fermi chopper / velocity selector





Velocity selector

- Discriminate beam in *velocity / wavelength*
- $\Delta\lambda/\lambda \sim 10\%$




Fermi Choppers

- Discriminates beam in *time* and *wavelength* simultaneously





Crystal monochromators (and analyzers)

- Discriminate beam in *wavelength* by Bragg's law
- $\Delta\lambda/\lambda \sim 1\%$ (plus multiples $\lambda/2, \lambda/3, \dots$)




Beamline design

- By a suitable selection of: choppers, velocity selectors etc etc the neutron scatterer is able to 'design' the beam optimal for his/her sample.



Some samples studied


- Numerous, cross-disciplinary
- Materials in different states, eg.
 - Crystals
 - Powders
 - Molecules in solution



- Material behaviour/function
 - Materials for fuel cells, batteries...
 - Magnets
 - Superconductors
 - Chemical reactions
 - Protein folding
 - Polymers
 - Metallurgy
 - ...




Introduction to McStas



- Å to m distances
- Fourier (reciprocal space) methods
- Direct space methods

Detectors

- Since neutrons are electrically neutral, they are difficult to detect.
- The preferred reaction is:




due to the high cross-section
E field → protons collected → signal

- Recent years lack of ${}^3\text{He}$ has forced the community to look for alternatives:



due to the high neutron capture cross-section of ${}^{10}\text{B}$

The energetic nuclei ionize gas molecules which can be collected as signals




Monte Carlo techniques

- Los Alamos has since then developed and perfected many different monte carlo codes leading to what is today known as the codes MCNP5 and MCNPX
- State of the art is MCNPX (or soon the merged MCNP6 code) that features numerous (even exotic) particles
- MCNP was originally Monte Carlo Neutron Photon, later N-Particle
- Mainly used for high-energy particle descriptions in weapons, power reactors and routinely used for estimating dose rates and needed shielding
- Does not to date handle coherent scattering of neutrons due to the focus on high energies




Ray-tracing methods



- When neutrons move in “free space”, we use ray-tracing - but in most cases in direction source → detector
- Of course parabolas rather than straight lines are used to implement gravity

Elements of Monte-Carlo raytracing

- Instrument Monte Carlo methods implement coherent scattering effects
- Uses deterministic propagation where this can be done
- Uses Monte Carlo sampling of “complicated” distributions and stochastic processes and multiple outcomes with known probabilities are involved
 - i.e. inside scattering matter
- Uses the particle-wave duality of the neutron to switch back and forward between deterministic ray tracing and Monte Carlo approach



- Result: A realistic and efficient transport of neutrons in the thermal and cold range
- **McStas**: the code (of Risø origin) that encompass transport, beam-line and detector simulation, analysis framework

Neutron ray/package:

Weight (p): # neutrons (left) in the package


Coordinates (x,y,z)

Velocity (v_x, v_y, v_z)

Spin (s_x, s_y, s_z)

Time (t)










Instrument: positioning + transformation between sequential component coordinate systems, e.g. neutron source, crystal, detector.




What is McStas used for?

- Instrumentation
- Virtual experiments
- Data analysis
- Teaching

KU, DTU 2005-2012
INSIS, NIDS, ESS workshops





Example from ILL



Example from ILL – E.Farhi

How to get from MCNPX to McStas

- Based on the latest MCNPX ESS target station (bi-spectral) geometry from ESS-Bilbao we have developed a McStas component mimicking both geometry and spectra.
- We are also working on alternatives which transport the neutron state directly, thus avoiding loss of phase-space / making assumptions



Danish in-kind contributions

- 1.0 M€. Proton beam control (Søren Pape Møller, AU)
- 1.5 M€. Data Management and Software Center (Stig Skelboe, KU)
- 1.0 M€. Instrument simulation central office (Kim Lefmann KU, P.Willendrup DTU)
- 0.1 M€. Integrating moderator- and instrument simulations
(B.Lauritzen, P.Willendrup, E.Nonbøl, E.Klinkby DTU)
- 0.2 M€. Radio-ecology baseline (Mikael Jensen, Sven Nielsen DTU)
- 0.1 M€. MANTID – DMSC (Stig Skelboe, KU)
- 0.8 M€. 5 DK-CH instrument packages
(Niels Bech Christensen, DTU; Christian Rüegg, PSI)



EUROPEAN
SPALLATION
SOURCE




Data Management and Software

- Staff: Stig Skelboe, Thomas Rod, Lars Melvyn, (secretary) (3 FTE)
- Supporters: scientists from KU, DTU-Risø, AU
- DMSC scope (under planning)
 - User service
 - Instrument control
 - Data acquisition
 - Data archiving
 - Data visualization and analysis (science modeling)
 - Instrument simulation




Instrument simulations




- Simulate a suite of simple instruments to investigate time structure
 - Later: move towards detailed instrument descriptions
- Answer questions from ESS instrument responsables
 - Compare thermal powder diffraction designs
 - Compare thermal spectrometer designs
 - Analyze effect of off-specular scattering
- Prepare for virtual experiments for data analysis
 - Event mode data; bootstrap
 - Effect of the pulse tail
 - Effect of multiple scattering; sample environment
- Maintain and develop McStas
- Study guide systems
 - Long thermal guides
 - Guide bundles
 - Bi-spectral extraction
- Support function for simulators



Example: Cold chopper spectrometer



- 100 m elliptical guide
- Wavelength multiplication at sample
- 30 (300) times IN5 flux
- Count rates of the order 10^8 / sec.
- VE shows expected resolution
- Moderator "Hot spot" is highly beneficial



Conclusions

- Danish universities are/will be heavy involved in many aspects of the design, construction and usage of the ESS and its instruments, including:
 - Data management
 - Radioecology
 - Instrument simulation
 - Instrument design
 - Neutron scattering experiments
 - Neutronics
 - Develop/maintain McStas




EUROPEAN
SPALLATION
SOURCE




Backup slides...

Overview



Moderators... (Where McStas starts)





Per beamline:

$$I_{BL}(x,y,E,t) = \frac{\Omega_{BL}(x,y)}{4\pi} I(x,y,E,t)$$



Example suite: 7 TOF spectrometers:

- ESS_IN5_reprate.instr
- ILL_BRISP.instr (Small-angle)
- ILL_H15_IN6.instr
- ILL_H16_IN5.instr
- ISIS_Hetfull.instr
- PSI_Focus.instr
- templateTOF.instr




Example suite: 5 TAS



- ILL_H142_IN12.instr
- ILL_H25_IN22.instr
- h8_test.instr
- templateTAS.instr
- linup-1.instr (Risø TAS 1)
- linup-2.instr
- linup-3.instr
- linup-4.instr
- linup-5.instr
- linup-6.instr
- linup-7.instr





Example suite: 1 Hybrid spectrometer + 1 Spin-echo



Spin-echo B scan dependence of wavelength






Example suite: Large scale structures



Example suite:

Diffractometers

- ILL_D1A.instr
- PSI_DMC.instr
- templateDIFF.instr
- templateLaue.instr



Example suite: Imaging

